13

UTILIZATION OF DRIED BAKERY PRODUCTS AS SUPPLEMENT IN BROILER RATION.

THESIS

PABLO R. DE LUNAS JR.

College of Agriculture, Forestry, Environment and Natural Resources CAVITE STATE UNIVERSITY Indang, Cavite

A 57 1 3003

UTILIZATION OF DRIED-BAKERY PRODUCTS AS SUPPLEMENT IN BROILER RATION

Undergraduate Thesis Submitted to the Faculty of the Cavite State University Indang, Cavite

In partial fulfillment of the requirements for the degree of Bachelor of Science in Agriculture (Major in Animal Science)



Utilization of dried bakery products a supplement in broiler ration 636.513 D38 2003

PABLO R. DE LUNAS JR.

April 2003

ABSTRACT

DE LUNAS, PABLO, JR. R. Utilization of Dried-Bakery Products as Supplement in Broiler Ration. Undergraduate Thesis. Bachelor of Science in Agriculture major in Animal Science (Animal Nutrition). Cavite State University, Indang, Cavite. April 2003. Adviser: Prof. Eraño C. Esguerra.

A study was conducted at the experimental area of the Animal Broiler Project of the Office of the Business Affairs (OBA) from November 2002 to December 2002 to: determine the effect of dried-bakery products in terms of growth rate, feed consumption and feed conversion efficiency of broiler; determine the optimum level of dried-bakery products for broiler ration; and determine the economics of incorporating dried-bakery products to commercial feeds.

Four treatment groups with a total of 100 day old chicks were used with different level of supplementation as follows: Treatment I (pure commercial feeds), Treatment II (90% commercial feeds + 10% dried-bakery products), Treatment III (85% commercial feeds + 15% dried-bakery products); and Treatment IV (80% commercial feeds + 20% dried-bakery products).

The results were insignificantly different (P>0.05) in final weight, final gain in weight and feed conversion efficiency of the broiler chicken. Although feed consumption were significantly different (P<0.01), birds with 20% dried-bakery products had a much higher feed consumption over the other groups.

Mortality rate was very low and occurred only in Treatment IV-replication 2, at the second week of age of rearing period.

Treatment IV was more profitable than the other treatment groups.

TABLE OF CONTENTS

	Page
BIOGRAPHICAL DATA	iii
ACKNOWLEDGMENT	iv
ABSTRACT	vi
TABLE OF CONTENTS	vii
LIST OF TABLES	ix
LIST OF APPENDIX TABLES	. x
LIST OF APPENDIX FIGURES	. xi
INTRODUCTION	. 1
Importance of the Study	. 2
Objectives of the Study	. 2
Limitation of the Study	. 3
Time and Place of the Study	. 3
DEFINITION OF TERMS	
REVIEW OF RELATED LITERATURE	
METHODOLOGY	. 10
Materials	. 11
Experimental Design and Treatments	. 11
Preparation of Housing	. 11
Preparation of Dried-Bakery Products	. 11
Feeding and Watering	. 12

Care and Management	12		
Health and Sanitation	12		
Marketing	13		
Data gathering	13		
Statistical Analysis	13		
RESULT AND DISCUSSION	14		
Body Weight	14		
Initial Feed Consumption, Weekly Cumulative Feed Consumption and Feed Efficiency of the Birds	15		
Cost and Return Analysis	17		
SUMMARY, CONCLUSIONS AND RECOMMENDATIONS			
Summary	19		
Conclusions	20		
Recommendations	20		
BIBLIOGRAPHY	21		
APPENDICES			

LIST OF TABLES

Table		Page
1	Initial weight gain in weight, and final body weight of the bird	14
2	Initial feed consumption, weekly cumulative feed consumption and feed efficiency	16
3	Cost and return analysis	18

LIST OF APPENDIX TABLES

Appendix Table		Page
1	Proximate Analysis	24
2	Analysis of variance for initial body weight of the birds	25
3	Analysis of variance for body weight one week after rearing	25
4	Analysis of variance for body weight two weeks after rearing	25
5	Analysis of variance for body weight three weeks after rearing	26
6	Analysis of variance for final body weight	26
7	Analysis of variance for feed consumption three weeks of age	26
8	Analysis of variance for feed consumption four weeks of age	. 27
9	Analysis of variance for feed consumption five weeks of age	. 27
10	Analysis of variance for feed consumption six weeks of age	. 27
11	Analysis of variance for feed conversion efficiency	. 28

LIST OF APPENDIX FIGURES

Appendix Figure		Page
1	Bakery shop in Balayan, Batangas	30
2	The dried-bakery product subjected to sun drying	31
3	The dried-bakery product subjected to carajay and fire wood for drying	32
4	The author pulverizing dried-bakery products using mortar and pestle	33
5	Process of separating fine from coarse particles of dried-bakery products	34
6	The finished bakery products ready for mixing with commercial feeds	35
7	The broiler house used in the study	36
8	The growing cages used in the study	37
9	Birds inside the cages at 2weeks of age	38
10	The author mixing the dried-bakery products with commercial feeds	39
11	Birds at 3 weeks of age	40
12	Birds at 4 weeks of age	41

UTILIZATION OF DRIED-BAKERY PRODUCTS AS SUPPLEMENT IN BROILER RATION $^{1\!\!/}$

Pablo Ramos de Lunas Jr.

LA thesis manuscript presented to the faculty of Department of Animal Science, College of Agriculture, Forestry, Environment and Natural Resources Cavite State University, Indang, Cavite in partial fulfillment of the requirements for the degree of Bachelor of Science in Agriculture major in Animal Science with Contribution No. BSA-2003-01-108. Prepared under the supervision of Prof. Eraño C. Esguerra.

INTRODUCTION

The size of poultry enterprise ranges from small farm flocks to large commercial operators. However, there has been an increased interest in recent years in keeping small flocks for the production of eggs or meat for home use. Regardless of the size of the enterprise, there are three important factors for the success of poultry raising: proper feeding, good management, and sanitation.

Broiler production operations involve raising chickens for meat. High quality rations are fed to secure rapid efficient gains. Generally, several flocks of birds are fed and marketed each year. Replacement pullet operations involve raising chicks, which are then sold, to egg production, or broiler production operation, depending on the type of bird raised (Gillespie, 1997).

Dried-bakery products is a feed produced form reclaimed (unused or stale) bakery-products or other materials such as candy, inedible flours, unusable nuts, and the like. The materials are blended and ground to produce a feedstuff that contains 9% to 12%