PLORAL BIODIVERSITY ASSESSMENT OF TIMALAN RIVER WATERSHED, CAVITE, PHILIPPINES

Undergraduate Thesis
Submitted to the Faculty of the
College of Agriculture, Food, Environment, and Natural Resources
Cavite State University
Indang, Cavite

In partial fulfillment
of the requirements for the degree
Bachelor of Science in Environmental Science



Floral biodiversity assessment of Timalan River Watershed, Cavite Philippines 333.95 C88 2018

CYJANE U. ALCAZAR ELIZA JOYCE T. CRUZ June 2018

ABSTRACT

ALCAZAR, CYJANE U. and CRUZ, ELIZA JOYCE T. Floral Biodiversity Assessment of Timalan River Watershed, Cavite, Philippines. Undergraduate thesis. Bachelor of Science in Environmental Science. Cavite State University, Indang, Cavite. June 2017. Adviser For. Junser P. Magpantay.

The study was conducted in Timalan River Watershed from August 2017 to April 2018. Specifically, this study aims to: (1) classify the land use of Timalan River Watershed (TRW); (2) identify the floral species that are found in TRW; (3) determine the number of floral species that can be found in TRW; (4) analyze the distribution of floral species within TRW; and (5) identify the indigenous floral species present in TRW. The data were gathered through ocular visits, actual field survey, literature review and secondary data. A land use map obtained from DENR-PENRO-Cavite (2017) was utilized to identify the land use types in TRW. Data collection and identification of floral species were conducted on ten sites in grassland, ten sites in forest, ten sites in mixed orchard, and ten sites in agricultural areas, which give a total of 40 selected sampling sites in TRW. Quadrat sampling method was done for every site.

Timalan River Watershed has a total area of 2,557.53 ha of grassland, 14.31 ha of forest, 0.36 ha of mixed orchard and 1,121.94 ha of agricultural areas. Grassland has the highest total area among the other land use types. A total of 92 floral species were identified and categorized into trees, intermediate and understorey. The indigenous floral species have almost the same number of exotic species in TRW.

The results showed that the average Shannon-Wiener Diversity Index of the whole TRW is low to very low, which ranged from 1.14 to 2.57. These results indicate that TRW has low diversity because there are relatively few species in the area. The

average Shannon's Evenness Index score is high to very high, which ranged from 1.55 to 1.00 due to having equal number of frequencies.

The indigenous floral species in Timalan River Watershed are threatened and slowly dominated by exotic species like mahogany. There is a need to conserve the indigenous floral species of TRW through programs like tree planting which will benefit the environment and the community.

TABLE OF CONTENTS

	Page
TITLE PAGE	i
APPROVAL SHEET	ii
BIOGRAPHICAL DATA	iii
ACKNOWLEDGEMENT. ABSTRACT.	v
LIST OF TABLES	xi
LIST OF FIGURES	xiv
LIST OF APPENDIX TABLES	xvi
LIST OF APPENDIX FIGURES	xviii
LIST OF APPENDICES	XX
INTRODUCTION	1
Statement of the Problem	3
Objectives of the Study	3
Significance of the Study	4
Time and Place of the Study	4
Scope and Limitations of the Study	5
Definition of Terms	5
Conceptual Framework of the Study	7
REVIEW OF RELATED LITERATURE	8
METHODOLOGY	15

	Research Design	15
	Materials	16
	Methods	16
	Study area	16
	Validation of land use map	19
	Ocular visit	19
	Sampling sites identification	19
	Data collection and identification of floral species	23
	Data analysis and interpretation	24
RESU	LTS AND DISCUSSION	27
	Land Use Map of Timalan River Watershed	27
	Floral Species That Can Be Found in Timalan River Watershed	36
	Occurrence of Plant Species According to Land Use Types	71
	Distribution of Floral Species in Timalan River Watershed using Shannon-Wiener and Shannon's Evenness	82
	Importance Value Index	88
	Indigenous Floral Species Found in Timalan River Watershed	91
SUMN	MARY, CONCLUSION, AND RECOMMENDATIONS	98
	Summary	98
	Conclusion	101
	Decommendation	103

REFERENCES	105
APPENDIX TABLES	107
APPENDIX FIGURES	141
APPENDICES	163